

Effect of Body Mass Index on Left Ventricular Mass in Career Male Firefighters



Maria Korre, ScD^{a,b}, Luiz Guilherme G. Porto, PhD^{a,c}, Andrea Farioli, MD^{a,b,d}, Justin Yang, MD, MPH^a, David C. Christiani, MD, MPH^a, Costas A. Christophi, PhD^{a,e}, David A. Lombardi, PhD^{a,f}, Richard J. Kovacs, MD^g, Ronald Mastouri, MD^g, Siddique Abbasi, MD^h, Michael Steigener, MDⁱ, Steven Moffatt, MD^j, Denise Smith, PhD^k, and Stefanos N. Kales, MD, MPH^{a,b,*}

Left ventricular (LV) mass is a strong predictor of cardiovascular disease (CVD) events; increased LV mass is common among US firefighters and plays a major role in firefighter sudden cardiac death. We aim to identify significant predictors of LV mass among firefighters. Cross-sectional study of 400 career male firefighters selected by an enriched randomization strategy. Weighted analyses were performed based on the total number of risk factors per subject with inverse probability weighting. LV mass was assessed by echocardiography (ECHO) and cardiac magnetic resonance, and normalized (indexed) for height. CVD risk parameters included vital signs at rest, body mass index (BMI)—defined obesity, obstructive sleep apnea risk, low cardiorespiratory fitness, and physical activity. Linear regression models were performed. In multivariate analyses, BMI was the only consistent significant independent predictor of LV mass indexes (all, $p < 0.001$). A 1-unit decrease in BMI was associated with 1-unit ($\text{g}/\text{m}^{1.7}$) reduction of LV mass/height^{1.7} after adjustment for age, obstructive sleep apnea risk, and cardiorespiratory fitness. In conclusion, after height-indexing ECHO-measured and cardiac magnetic resonance—measured LV mass, BMI was found to be a major driver of LV mass among firefighters. Our findings taken together with previous research suggest that reducing obesity will improve CVD risk profiles and decrease on-duty CVD and sudden cardiac death events in the fire service. Our results may also support targeted noninvasive screening for LV hypertrophy with ECHO among obese firefighters. © 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). (Am J Cardiol 2016;118:1769–1773)

Despite the critical prognostic significance of left ventricular (LV) mass,^{1–3} its measurement and role in clinical practice have yet to be established.⁴ Echocardiography (ECHO) and cardiac magnetic resonance (CMR) are the 2 most commonly used imaging methods for the assessment of LV mass. Although, CMR is considered the gold standard for assessing LV mass, ECHO is a well validated, noninvasive method that is more widely used in clinical practice.⁵ In addition to considering different imaging methods, disagreement exists as to the most appropriate method of indexing LV mass to body size parameters.³ Current evidence suggests indexing by height to the allometric powers

of 1.7 and 2.7 are the most accurate normalization techniques.^{4,6,7} This study identifies the most important predictors of LV mass after indexing for height among career male firefighters as assessed by both ECHO and CMR.

Methods

Male career firefighters, aged 18 years and older were recruited from the Indianapolis Fire Department. Eligible firefighters had a recorded fire department—sponsored medical examination in the last 2 years that included a submaximal exercise tolerance test and had no restrictions on duty.

^aDepartment of Environmental Health, Environmental and Occupational Medicine and Epidemiology Program, Harvard T.H. Chan School of Public Health, Boston, Massachusetts; ^bDepartment of Occupational Medicine, Cambridge Health Alliance, Harvard Medical School, Cambridge, Massachusetts; ^cFaculty of Physical Education, Cardiovascular Laboratory of the Faculty of Medicine, University of Brasilia, Brasilia, Brazil; ^dDepartment of Medical and Surgical Sciences, University of Bologna, Italy; ^eCyprus International Institute for Environmental and Public Health, Cyprus University of Technology, Limassol, Cyprus; ^fCenter for Injury Epidemiology, Liberty Mutual Research Institute for Safety, Hopkinton, Massachusetts; ^gDepartment of Medicine, Krannert Institute of Cardiology, Indiana University, Indianapolis, Indiana; ^hDepartment of Cardiology, Providence Veteran Affairs Medical Center, Cardiovascular Institute of the Alpert Medical School of Brown University, Providence, Rhode Island; ⁱHeart &

Vascular Center, Brigham and Women's Hospital, Boston, Massachusetts; ^jPublic Safety Medical, Indianapolis, Indiana; and ^kDepartment of Health and Exercise Sciences, Skidmore College, Saratoga Springs, New York, New York. Manuscript received June 7, 2016; revised manuscript received and accepted August 14, 2016.

This investigation was supported by the Federal Emergency Management Agency (FEMA) Assistance to Firefighters Grant (AFG) program's award EMW-2011-FP-00663 (PI: Dr. Kales). LGGP is a Harvard School of Public Health Visiting Scientist supported by PDE: nr 207136/2014.9 from the Conselho Nacional de Desenvolvimento Científico e Tecnológico—CNPq—Brazil.

See page 1772 for disclosure information.

*Corresponding author: Tel: (617) 665-1580; fax: (617) 665-1672.

E-mail address: skales@hsph.harvard.edu (S.N. Kales).

Table 1
Baseline descriptive characteristics

Variables	Study Sample (N = 393)	Study Sample Unweighted
Age (years)*	47 ± 8.2	45 ± 8.1
Height (cm)*	179 ± 6.4	179 ± 6.6
Heart rate (bpm)*	81 ± 13	80 ± 13
Body weight (kg)*	99 ± 17	97 ± 17
Resting systolic blood pressure (mm Hg)*	126 ± 9.7	125 ± 9.4
Resting diastolic blood pressure (mm Hg)*	82 ± 8.1	81 ± 7.4
High risk of obstructive sleep apnea†	112 (38%)	254 (32%)
Body mass index (kg/m ²)*	31 ± 4.6	30 ± 4.5
Smoker†	50 (13%)	135 (13%)
History of heart rhythm problems†	60 (16%)	153 (15%)
Family history of cardiac problems†	153 (40%)	426 (41%)
Age ≥ 40 (years)†	301 (78%)	770 (73%)
Body mass index ≥ 30 (kg/m ²)†	260 (56%)	474 (45%)
Low cardiorespiratory fitness†	178 (47%)	363 (34%)
Moderate to vigorous physical activity (min/week)*	177 ± 117	187 ± 118
LV mass by echocardiography (g)*	189 ± 38	187 ± 37
LV mass by cardiac magnetic resonance (g)*	139 ± 24	138 ± 23
LV mass by echocardiography indexed to height ^{1.7} (g/m ^{1.7})*	70 ± 13	70 ± 13
LV mass by echocardiography indexed to height ^{2.7} (g/m ^{2.7})*	40 ± 7.6	39 ± 7.4
LV mass by cardiac magnetic resonance indexed to height ^{1.7} (g/m ^{1.7})*	52 ± 8.3	51 ± 8.1
LV mass by cardiac magnetic resonance indexed to height ^{2.7} (g/m ^{2.7})*	29 ± 4.7	29 ± 4.6

bpm = beats per minute; LV = left ventricular.

* Mean (SD) for continuous variables.

† n (%) for categorical variables.

From those eligible, we selected a total of 400 participants, using an “enriched” randomization strategy based on age at randomization, obesity, hypertension, and cardiorespiratory fitness status at last examination, where a larger number of higher risk participants could be selected. Thus, we randomly selected 100 participants from the entire eligible population; 75 low-risk participants (age <40, nonobese, free of hypertension, and high cardiorespiratory fitness) and 225 higher risk participants (at least 2 of the following: age ≥40 years, obese, hypertension, or low cardiorespiratory fitness) for further LV hypertrophy/cardiomegaly screening and imaging tests. Obesity was defined by standard criteria (body mass index [BMI] ≥30 kg/m²). Hypertension was considered present if blood pressure at rest is ≥140/90 mm Hg. Low cardiorespiratory fitness was defined as the bottom tertile, as measured by the recorded treadmill time and the estimated maximal VO₂ during the last exercise test. Those selected were included in the study if they had no contraindication to CMR and signed informed consent to participate.

LV mass was assessed by both ECHO and CMR imaging. First, a transthoracic cardiac echocardiogram was done as a simple 2-dimensional study with limited m-mode recordings. An abbreviated CMR was performed as “function only” immediately after the ECHO. Images were obtained using a retrospectively electrocardiogram gated steady-state free precession cine sequence. In this fashion, a contiguous short-axis stack of 8-mm slices was obtained parallel to the atrioventricular groove to cover the entire length of the LV. Then, manual tracing of end-diastolic epicardial and endocardial borders was performed. Standard long-axis views were also obtained including horizontal long-axis, vertical

long-axis, and 3-chamber views, facilitating the interpretation of ventricular function. Board certified specialists performed clinical interpretation of imaging. LV mass indexes were derived by dividing LV mass in grams with height to the allometric powers of 1.7 and 2.7 (in meters^{1.7} and meters^{2.7}, respectively).

Height was measured in the standing position with a clinic stadiometer. Body weight was measured with bare feet and in light clothes on a calibrated scale. BMI was calculated as the weight in kilograms divided by the square of height in meters. Blood pressure was measured using an appropriately sized cuff with the subject in the seated position. Heart rate and blood pressure were obtained in a resting state from the physical examination (and were not measured before the exercise test). Medical examination data were further supplemented by a preimaging questionnaire, which collected comprehensive information on smoking status, a history of heart rhythm problems, family history of cardiac problems, and moderate to vigorous physical activity level in minutes per week. Obstructive sleep apnea risk was assessed using the validated Berlin Questionnaire.⁸

We performed a weighted analysis so as to account for our enriched randomization sampling strategy. Weights were calculated based on the total number of risk factors per subject with the technique of inverse probability weighting. Baseline characteristics were described using the mean (SD) in the case of quantitative variables and the frequency (%) for categorical variables. The effects of the different independent variables on the LV mass indexes were assessed with the use of linear regression models. Any independent variables that were significant in the univariate regression models were included in

Table 2

Simple linear regression models of cardiovascular risk factors and LV mass assessed by echocardiography and cardiac magnetic resonance and normalized for height to allometric powers of 1.7 and 2.7 as continuous variable

Variables	Assessed by ECHO				Assessed by CMR			
	Model 1*		Model 2 [†]		Model 3*		Model 4 [†]	
	β (SE)	P	β (SE)	p	β (SE)	p	β (SE)	p
Age (years)	0.11 (0.1)	0.02 [‡]	0.10 (0.0)	<0.01 [‡]	0.01 (0.1)	0.80	0.03 (0.0)	0.18
Heart Rate (bpm)	-0.01 (0.03)	0.78	-0.02 (0.0)	0.33	-0.02 (0.0)	0.34	-0.02 (0.0)	0.08
Resting systolic blood pressure (mmHg)	0.13 (0.04)	<0.01 [‡]	0.08 (0.0)	<0.01 [‡]	0.17 (0.0)	<0.01 [‡]	0.10 (0.0)	<0.01 [‡]
Resting diastolic blood pressure (mmHg)	-0.002 (0.1)	0.97	0.002 (0.0)	0.96	0.14 (0.1)	<0.01 [‡]	0.08 (0.0)	<0.01 [‡]
High risk of obstructive sleep apnea	5.64 (0.99)	<0.01 [‡]	3.12 (0.6)	<0.01 [‡]	3.90 (0.6)	<0.01 [‡]	2.20 (0.3)	<0.01 [‡]
Body Mass Index (kg/m ²)	0.95 (0.1)	<0.01 [‡]	0.51 (0.1)	<0.01 [‡]	0.86 (0.1)	<0.01 [‡]	0.46 (0.0)	<0.01 [‡]
Smoker	-2.62 (1.2)	0.03 [‡]	-1.3 (0.7)	0.05 [‡]	-1.11 (0.77)	0.15	-0.63 (0.4)	0.15
History of Heart Rhythm Problems	-2.68 (1.2)	0.02 [‡]	-1.48 (0.6)	0.02 [‡]	-0.98 (0.75)	0.19	-0.57 (0.4)	0.19
Family History of cardiac problems	3.26 (0.8)	<0.01 [‡]	2.01 (0.5)	<0.01 [‡]	0.07 (0.6)	0.89	0.24 (0.5)	0.44
Low cardiorespiratory fitness	3.24 (0.8)	<0.01 [‡]	1.94 (0.5)	<0.01 [‡]	1.53 (0.6)	<0.01 [‡]	0.95 (0.3)	<0.01 [‡]
Moderate to vigorous Physical Activity (min/week)	-0.02 (0.0)	<0.01 [‡]	-0.01 (0.0)	<0.01 [‡]	-0.002(0.0)	0.46	-0.001 (0.0)	0.28

bpm = beats per minute; CMR = cardiac magnetic resonance; ECHO = echocardiography.

* LV mass normalized for height to the allometric power of 1.7.

[†] LV mass normalized for height to the allometric power of 2.7.

[‡] Statistically significant p-values.

the multivariate regression models. In the multivariate analysis, we followed the backward stepwise elimination process with a removal criterion of $\alpha = 0.20$. Then, considering the predictors that resulted from the backward elimination process and variables that we knew a priori to be important clinical predictors, we constructed the final multivariate regression models. The interaction effects between BMI with obstructive sleep apnea risk and age were also assessed in these models. Collinearity was evaluated using the variance inflation factor. Analyses were performed using SPSS, version 21.0 (IBM, Armonk, New York). A p value of <0.05 was considered statistically significant, and all tests performed were 2 sided.

Results

Of the 400 firefighters, we excluded 7 participants with missing measurements of LV mass, assessed by CMR. Baseline characteristics are summarized in Table 1. The mean age of the study subjects was 45 (8.1) years, their mean BMI was 30 (4.5) kg/m² and 45% were obese.

The univariate analyses summarized in Table 2 revealed highly statistically significant associations between both LV mass height indexes, assessed by both ECHO and CMR, with systolic blood pressure at rest, hypertension, high risk of obstructive sleep apnea, low cardiorespiratory fitness, and BMI (all $p < 0.01$). Age, family history of cardiac problems, and physical activity also showed a significant association with both LV mass indexes, when LV mass was based on ECHO measurement (at least $p < 0.01$).

In all 4 models evaluated, namely with LV mass assessed by ECHO or CMR and normalized with height to either 1.7 or 2.7, only BMI was consistently associated with LV mass in a statistically significant fashion ($p < 0.001$) in all multivariate models.

Final multivariate regression models showing the associations between the statistically and clinically significant

predictors of LV mass are summarized in Table 3. A 1-unit decrease in BMI was associated with 1 unit (g/m^{1.7}) reduction of LV mass/height^{1.7} after adjustment for age, obstructive sleep apnea risk, and cardiorespiratory fitness.

Discussion

The present cross-sectional study in US firefighters using ECHO and CMR measurements found BMI to be the strongest and most consistent independent predictor of LV mass indexed by height. In simple linear regression models, apart from BMI, the associations were highly statistically significant for high risk of obstructive sleep apnea, systolic blood pressure at rest, and low cardiorespiratory fitness consistently in all 4 models ($p < 0.01$). In multivariate models, however, BMI was the only consistently significant predictor. Therefore, our study clearly supported BMI as a major determinant of LV mass.

Given the epidemic level of obesity in the US fire service, it is not surprising that we found BMI to be the strongest predictor of LV mass in this population. This is consistent with the studies, which find obesity to be a risk factor for LV hypertrophy and increased cardiac mass.^{9,10} In addition, given that obesity is associated with cardiovascular disease (CVD) risk factor clustering,^{11,12} it probably explains why other factors such as blood pressure and obstructive sleep apnea risk were weaker predictors because of their association with LV mass may be closely linked to their association or co-morbidity with obesity.¹³ Given our previous findings that obesity-associated sudden cardiac death (SCD) among younger firefighters was largely driven by an increased cardiac mass in SCD victims compared with controls,¹⁴ our results reinforce that decreasing obesity in the fire service will lower the risk of LV hypertrophy and on-duty CVD events, particularly SCD. In agreement with findings that even small reductions on BMI may produce significant beneficial effects on metabolic syndrome and

Table 3

Multivariate linear regression models of cardiovascular risk factors and LV mass assessed by echocardiography and cardiac magnetic resonance and normalized for height to allometric powers of 1.7 and 2.7 as continuous variable

R ²	Assessed by ECHO				Assessed by CMR			
	Model 1*		Model 2 [†]		Model 3*		Model 4 [†]	
	0.134		0.125		0.239		0.215	
	β (SE)	P	β (SE)	p	β (SE)	p	β (SE)	p
Age (years)	0.04 (0.1)	0.52	0.07 (0.0)	0.07	0.02 (0.0)	0.56	0.04 (0.0)	0.05
High risk of obstructive sleep apnea	0.76 (1.1)	0.49	0.36 (0.6)	0.57	0.15 (0.6)	0.81	0.11 (0.4)	0.76
Body Mass Index (kg/m ²)	1.01 (0.1)	<0.001 [‡]	0.55 (0.1)	<0.001 [‡]	0.83 (0.1)	<0.001 [‡]	0.45 (0.0)	<0.001 [‡]
Low cardiorespiratory fitness	-0.23 (1.0)	0.83	-0.34 (0.6)	0.57	-0.96 (0.6)	0.10	-0.70 (0.3)	0.05

CMR = cardiac magnetic resonance; ECHO = echocardiography.

* LV mass normalized for height to the allometric power of 1.7.

[†] LV mass normalized for height to the allometric power of 2.7.

[‡] Statistically significant p values.

other CVD risk factors,^{15,16} our results suggest that a 1-unit decrease in BMI will reduce LV mass index by 1 unit (g/m^{1.7}). Our results may also support targeted noninvasive screening for LV hypertrophy with ECHO among obese firefighters.

Based on the values of R² for our final multivariate regression models, we were able to explain 12.5% to 13.4% of the variability of LV mass indexed by height to the allometric powers of 1.7 and 2.7 based on ECHO assessments and 21.5% to 23.9% based on CMR assessments. We were able to explain 10% more of the LV mass variability with the CMR models compared with ECHO ones, irrespective of the indexation technique. This is likely explained by CMR measurements that are more standardized across techniques and institutions and less dependent on operator's skill and experience, acoustic window adequacy, and LV mass geometric assumptions.^{3,17}

Our study has some modest limitations. Because of its cross-sectional design, we can only demonstrate associations and not causation; however, the findings are consistent with past studies¹⁸ and are biologically plausible. In addition, because of the very small number of participating women firefighters in our study, only male participants were included in the present study.

Our study also has a number of important strengths. We were able to collect comprehensive data on CVD risk factors from both medical examinations and a screening questionnaire. The BMI was measured during medical examinations, which avoided self-reporting biases. Obstructive sleep apnea risk was assessed by the widely used and validated Berlin Questionnaire, which has high sensitivity and specificity (86% and 77%, respectively) and demonstrates a high yield in public safety occupations.^{8,19} Moreover, we used imaging results for LV mass by both ECHO and CMR. Another important strength of our study is that we normalized LV mass by height to 2 different allometric powers, which allowed us to perform a more holistic assessment of its potential predictors. Furthermore, our results were consistent among the imaging methods and the indexing methods, making our findings more robust. Finally, our sample had similar anthropometric characteristics and CVD risk factors

to those found in other epidemiologic studies of firefighters.^{11,12,20} Therefore, we believe that our results could be generalized to most male career firefighters.

Acknowledgment: The authors would like to thank all the participating firefighters and the Indianapolis Fire Department; the staff and clinical leadership of the clinics who examined the firefighters; Dr. Carol Jisseth Zárate Ardila, MD and Dr. Konstantina Sampani, MD who helped with the data entry.

Disclosures

The authors have no conflicts of interest to disclose.

1. Levy D, Garrison RJ, Savage DD, Kannel WB, Castelli WP. Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* 1990;322:1561–1566.
2. Haider AW, Larson MG, Benjamin EJ, Levy D. Increased left ventricular mass and hypertrophy are associated with increased risk for sudden death. *J Am Coll Cardiol* 1998;32:1454–1459.
3. Armstrong AC, Gidding S, Gjesdal O, Wu C, Bluemke DA, Lima JA. LV mass assessed by echocardiography and CMR, cardiovascular outcomes, and medical practice. *JACC Cardiovasc Imaging* 2012;5:837–848.
4. Gidding SS. Controversies in the assessment of left ventricular mass. *Hypertension* 2010;56:26–28.
5. Bottini PB, Carr AA, Prisant LM, Flickinger FW, Allison JD, Gottdiener JS. Magnetic resonance imaging compared to echocardiography to assess left ventricular mass in the hypertensive patient. *Am J Hypertens* 1995;8:221–228.
6. Chirinos JA, Segers P, De Buyzere ML, Kronmal RA, Raja MW, De Bacquer D, Claessens T, Gillebert TC, St John-Sutton M, Rietzschel ER. Left ventricular mass: allometric scaling, normative values, effect of obesity, and prognostic performance. *Hypertension* 2010;56:91–98.
7. Cuspidi C, Meani S, Negri F, Giudici V, Valerio C, Sala C, Zanchetti A, Mancia G. Indexation of left ventricular mass to body surface area and height to allometric power of 2.7: is the difference limited to obese hypertensives? *J Hum Hypertens* 2009;23:728–734.
8. Webber MP, Lee R, Soo J, Gustave J, Hall CB, Kelly K, Prezant D. Prevalence and incidence of high risk for obstructive sleep apnea in World Trade Center-exposed rescue/recovery workers. *Sleep Breath* 2011;15:283–294.
9. Soteriades ES, Targino MC, Talias MA, Hauser R, Kawachi I, Christiani DC, Kales SN. Obesity and risk of LVH and ECG abnormalities in US firefighters. *J Occup Environ Med* 2011;53:867–871.

10. Soteriades ES, Smith DL, Tsismenakis AJ, Baur DM, Kales SN. Cardiovascular disease in US firefighters: a systematic review. *Cardiol Rev* 2011;19:202–215.
11. Soteriades ES, Hauser R, Kawachi I, Liarokapis D, Christiani DC, Kales SN. Obesity and cardiovascular disease risk factors in firefighters: a prospective cohort study. *Obes Res* 2005;13:1756–1763.
12. Tsismenakis AJ, Christophi CA, Burrell JW, Kinney AM, Kim M, Kales SN. The obesity epidemic and future emergency responders. *Obesity (Silver Spring)* 2009;17:1648–1650.
13. Kales SN, Tsismenakis AJ, Zhang C, Soteriades ES. Blood pressure in firefighters, police officers, and other emergency responders. *Am J Hypertens* 2009;22:11–20.
14. Yang J, Teehan D, Farioli A, Baur DM, Smith D, Kales SN. Sudden cardiac death among firefighters ≤ 45 years of age in the United States. *Am J Cardiol* 2013;112:1962–1967.
15. Mileski KS, Leitao JL, Lofrano-Porto A, Grossi Porto LG. Health-related physical fitness in middle-aged men with and without metabolic syndrome. *J Sports Med Phys Fitness* 2015;55:223–230.
16. Elmer PJ, Obarzanek E, Vollmer WM, Simons-Morton D, Stevens VJ, Young DR, Lin PH, Champagne C, Harsha DW, Svetkey LP, Ard J, Brantley PJ, Proschan MA, Erlinger TP, Appel LJ. Effects of comprehensive lifestyle modification on diet, weight, physical fitness, and blood pressure control: 18-month results of a randomized trial. *Ann Intern Med* 2006;144:485–495.
17. Celebi AS, Yalcin H, Yalcin F. Current cardiac imaging techniques for detection of left ventricular mass. *Cardiovasc Ultrasound* 2010;8:19.
18. Mann CJ. Observational research methods. Research design II: cohort, cross sectional, and case-control studies. *Emerg Med J* 2003;20:54–60.
19. Baur DM, Christophi CA, Tsismenakis AJ, Cook EF, Kales SN. Cardiorespiratory fitness predicts cardiovascular risk profiles in career firefighters. *J Occup Environ Med* 2011;53:1155–1160.
20. Kales SN, Polyhronopoulos GN, Aldrich JM, Leitao EO, Christiani DC. Correlates of body mass index in hazardous materials firefighters. *J Occup Environ Med* 1999;41:589–595.